Author Index

ALLENDE, L.M., HERNÁNDEZ, M., CORELL, A., GARCÍA-PÉREZ, M.A., VARELA, P., MORENO, A., CARAGOL, I., GARCÍA-MARTÍN, F., GUILLÉN-PERALES, J., OLIVÉ, T., ESPAÑOL, T. & ARNAIZ-VILLENA, A. A novel CD18 genomic deletion in a patient with severe leucocyte adhesion deficiency: a possible CD2/lymphocyte function-associated antigen-1 functional association in humans, 440
ALTMANN, D.M. *see* DOUEK, D.C.
ALTMANN, K. *see* GILL, H.S.
ALTUWAIJRI, S. *see* THATCHER, T.H.
ANAND, A. *see* BODMAN-SMITH, M.D.
AONO, A., ENOMOTO, H., YOSHIDA, N., YOSHIZAKI, K., KISHIMOTO, T. & KOMORI, T. Forced expression of terminal deoxynucleotidyl transferase in fetal thymus resulted in a decrease in γδ T cells and random dissemination of Vγ3Vδ1 T cells in skin of newborn but not adult mice, 489
APPELBERG, R. *see* SILVA, R.A.
ARNAIZ-VILLENA, A. *see* ALLENDE, L.M.
AUSTYN, J. *see* BEHBOUDI, S.
AZORSA, D.O. *see* HIGGINBOTTOM, A.

BAIRD, M. *see* FAULKNER, L.
BANDYOPADHYAY, S. *see* BASU, A.
BANTING, G. *see* MAILE, R.
BAO, S., BEAGLEY, K.W., FRANCE, M.P., SHEN, J. & HUSBAND, A. J. Interferon-γ plays a critical role in intestinal immunity against *Salmonella typhimurium* infection, 464
BARAK, V. *see* HOURI-HADDAD, Y.
BARCENAS-MORALES, G., MERKENSCHLAGER, M., WAHID, F., DÖFFINGER, R. & IVANYI, J. Recessive expression of H2A-controlled immune response phenotype depends critically on antigen dose, 221
BARTH, R.K. *see* THATCHER, T.H.
BARTNES, K. & HANNESTAD, K. Native IgG2a^b is barely antigenic to major histocompatability complex class II-restricted T cells owing to inefficient internalization by professional antigen-presenting cells, 510
BASU, A., CHAKRABARTI, G., SAHA, A. & BANDYOPADHYAY, S. Modulation of CD11C$^+$ splenic dendritic cell functions in murine visceral leishmaniasis: correlation with parasite replication in the spleen, 305
BAUER, M. *see* KRANZER, K.
BEAGLEY, K.W. *see* BAO, S.
BEFUS, A.D. *see* DÉRY, R.E.
BEHBOUDI, S., CHAO, D., KLENERMAN, P. & AUSTYN, J. The effects of DNA containing CpG motif on dendritic cells, 361
BERGMEIER, L. *see* LEHNER, T.
BINGLEY, P.J. *see* KRETOWSKI, A.
BISSONNETTE, E.Y. *see* DÉRY, R.E.
BLUM, S. *see* WAGNER, C.
BODMAN-SMITH, M.D., ANAND, A., DURAND, V., YOUINOU, P.Y. & LYDYARD, P. M. Decreased expression of FcγRIII (CD16) by γδ T cells in patients with rheumatoid arthritis, 498
BRAVERY, C. *see* LEHNER, T.
BROMAGE, S.J. *see* SEARLE, R.F.
BROSH, N., EILAT, E., ZINGER, H. & MOZES, E. Characterization and role in experimental systemic lupus erythematosus of T-cell lines specific to peptides based on complementarity-determining region-1 and complementarity-determining region-3 of a pathogenic anti-DNA monoclonal antibody, 257
BUCHAN, G. *see* FAULKNER, L.
BUDDLE, B.M. *see* RHODES, S.G.
BÜRGER, A. *see* WAGNER, C.

CAMPBELL, S.B., KOMATA, T. & KELSO, A. Differential effects of CD4 and CD8 engagement on the development of cytokine profiles of murine CD4$^+$ and CD8$^+$ T lymphocytes, 394
CARAGOL, I. *see* ALLENDE, L.M.
CHAKRABARTI, G. *see* BASU, A.
CHAO, D. *see* BEHBOUDI, S.
CHIZZOLINI, C. *see* RIBBENS, C.
COLSTON, M.J. *see* TASCON, R.E.
CORELL, A. *see* ALLENDE, L.M.
CRANAGE, M. *see* LEHNER, T.
CROSS, M.L. *see* GILL, H.S.
CURRY, J.E. *see* SEARLE, R.F.

DALGLEISH, A.G. *see* TODRYK, S.M.
DAYAN, C.M. *see* MAILE, R.
DAYER, J.-M. *see* RIBBENS, C.
DENNIS, M. *see* LEHNER, T.
DÉRY, R.E., LIN, T.-J., BEFUS, A.D., MILNE, C.D., MOQBEL, R., MÉNARD, G. & BISSONNETTE, E. Y. Redundancy or cell-type-specific regulation? Tumour necrosis factor in alveolar macrophages and mast cells, 427
DÖFFINGER, R. *see* BARCENAS-MORALES, G.
DOUEK, D.C. & ALTMANN, D.M. T-cell apoptosis and differential human leucocyte antigen class II expression in human thymus, 249
DOYLE, C. *see* LEHNER, T.
DURAND, V. *see* BODMAN-SMITH, M.D.
DURRANT, S. *see* NICOL, A.

ECKES, B. *see* YAMAMOTO, T.
EGUCHI, K. *see* HIDA, A.
EILAT, E. *see* BROSH, N.
ELSEGOOD, K.A. *see* MAILE, R.
ELSON, C.J. *see* MACHT, L.M.
ENOMOTO, H. *see* AONO, A.
ESPAÑOL, T. *see* ALLENDE, L.M.
ESTES, D.M. *see* HAAS, K.M.
EVANS, W.H. *see* OVIEDO-ORTA, E.

FAULKNER, L., BUCHAN, G. & BAIRD, M. Interleukin-10 does not affect phagocytosis of particulate antigen by bone marrow-derived dendritic cells but does impair antigen presentation, 523
FISCHER, B. *see* MÜLLER, B.
FRANCE, M.P. *see* BAO, S.
FRANCIS, J.N., LAMONT, A.G. & THOMPSON, S. J. The route of administration of an immunodominant peptide derived from heat-shock protein 65 dramatically affects disease outcome in pristane-induced arthritis, 338
FUKADA, S. *see* IWAO, M.

GARCÍA-MARTÍN, F. *see* ALLENDE, L.M.
GARCÍA-PÉREZ, M.A. *see* ALLENDE, L.M.
GASTON, J.S.H. *see* MACHT, L.M.
GIBBS, B. *see* WELKER, P.
GILL, H.S., ALTMANN, K., CROSS, M.L. & HUSBAND, A. J. Induction of T helper 1- and T helper 2-type immune responses during *Haemonchus contortus* infection in sheep, 458
GILLESPIE, K.M. *see* KRETOWSKI, A.
GLUE, C. *see* JINQUAN, T.
GOMES, M.S. *see* SILVA, R.A.
GRABBE, J. *see* WELKER, P.
GRISWOLD, D.E. *see* PODOLIN, P.L.
GUILLÉN-PERALES, J. *see* ALLENDE, L.M.

HAAS, K.M. & ESTES, D. M. Activation of bovine B cells via surface immunoglobulin M cross-linking or CD40 ligation results in different B-cell phenotypes, 272
HALABI, A. *see* HOURI-HADDAD, Y.
HALL, G. *see* LEHNER, T.
HAMANO, S. *see* NINOMIYA, T.
HANNESTAD, K. *see* BARTNES, K.
HÄNSCH, G.M. *see* WAGNER, C.
HARADA, T. *see* IWAO, M.
HARDING, T.C. *see* MAILE, R.
HARTMANN, K. *see* YAMAMOTO, T.
HAYASHI, N. *see* KINGETSU, I.
HEEG, K. *see* KRANZER, K.
HENZ, B.M. *see* WELKER, P.
HERNÁNDEZ, M. *see* ALLENDE, L.M.
HEWISON, R.G. *see* RHODES, S.G.
HIDA, A., KAWAKAMI, A., NAKASHIMA, T., YAMASAKI, S., SAKAI, H., URAYAMA, S., IDA, H., NAKAMURA, H., MIGITA, K., KAWABE, Y. & EGUCHI, K. Nuclear factor-κB and caspases co-operatively regulate the activation and apoptosis of human macrophages, 553
HIGGINBOTTOM, A., WILKINSON, I., MCCULLOUGH, B., LANZA, F., AZORSA, D.O., PARTRIDGE, L.J. & MONK, P. N. Antibody cross-linking of human CD9 and the high-affinity immunoglobulin E receptor stimulates secretion from transfected rat basophilic leukaemia cells, 546
HIMENO, K. *see* SAKAI, T.
HIRAMINE, C. *see* IWAO, M.
HIRASAWA, N. *see* SHIRAISHI, M.
HIRST, E.M.A. *see* TASCON, R.E.
HISAEDA, H. *see* SAKAI, T.
HONJO, N. *see* LEE, M.-H.
HOURI-HADDAD, Y., SOSKOLNE, W.A., HALABI, A., BARAK, V. & SHAPIRA, L. Repeat bacterial challenge in a subcutaneous chamber model results in augmented tumour necrosis factor-α and interferon-γ response, and suppression of interleukin-10, 215
HOY, T. *see* OVIEDO-ORTA, E.
HUG, F. *see* WAGNER, C.
HUSBAND, A.J. *see* BAO, S.
see also GILL, H.S.

IDA, H. *see* HIDA, A.
IDE, T. *see* OHNO, N.
INOUYE, S. *see* KINGETSU, I.
see also OHNO, N.; TODA, M.
ISAJI, S. *see* KONDO, A.
ISHII, K. *see* SAKAI, T.
ISHIKAWA, H. *see* SAKAI, T.
IVANYI, J. *see* BARCENAS-MORALES, G.
IWAO, M., FUKADA, S., HARADA, T., TSUJIKAWA, K., YAGITA, H., HIRAMINE, C., MIYAGOE, Y., TAKEDA, S. & YAMAMOTO, H. Interaction of merosin (laminin 2) with very late antigen-6 is necessary for the survival of $CD4^+$ $CD8^+$ immature thermocytes, 481

JACOBI, H.H. *see* JINQUAN, T.
JINQUAN, T., QUAN, S., JACOBI, H.H., MADSEN, H.O., GLUE, C., SKOV, P.S., MALLING, H.-J. & POULSEN, L.K. CXC chemokine receptor 4 expression and stromal cell-derived factor-1α-induced chemotaxis in $CD4^+$ T lymphocytes are regulated by interleukin-4 and interleukin-10, 402
JUJI, T. *see* NICOL, A.
JUNT, T. *see* OEHEN, S.

KATAOKA, T. *see* LEE, M.-H.
KAWABE, Y. *see* HIDA, A.
KAWAKAMI, A. *see* HIDA, A.
KELSO, A. *see* CAMPBELL, S.B.
KEZUKA, T. & STREILEIN, J. W. Evidence for multiple CD95–CD95 ligand interactions in anterior chamber-associated immune deviation induced by soluble protein antigen, 451
KIMURA, G. *see* NINOMIYA, T.
KINALSKA, I. *see* KRĘTOWSKI, A.
KINGETSU, I., OHNO, N., HAYASHI, N., SAKAGUCHI, M., INOUYE, S. & SAITO, S. Common antigenicity between Japanese cedar (*Cryptomeria japonica*) pollen and Japanese cypress (*Chamaecyparis obtusa*) pollen, I. H-2 complex affects cross responsiveness to Cry j 1 and Cha o 1 at the T- and B-cell level in mice, 625
KIRWAN, J.R. *see* MACHT, L.M.
KISHIMOTO, T. *see* AONO, A.
KITAZAWA, T. & STREILEIN, J. W. Studies on delayed systemic effects of ultraviolet B radiation on the induction of contact hypersensitivity, 3. Dendritic cells from secondary lymphoid organs are deficient in interleukin-12 production and capacity to promote activation and differentiation of T helper type 1 cells, 296
KLENERMAN, P. *see* BEHBOUDI, S.
KOBAYASHI, Y. *see* SHIRAISHI, M.
KOEZUKA, Y. *see* NICOL, A.
KOMATA, T. *see* CAMPBELL, S.B.
KOMORI, T. *see* AONO, A.
KONDO, A., ISAJI, S., NISHIMURA, Y. & TANAKA, T. Transcriptional and post-transcriptional regulation of monocyte chemoattractant protein-3 gene expression in human endothelial cells by phorbol ester and cAMP signalling, 561
KRĘTOWSKI, A., GILLESPIE, K.M., BINGLEY, P.J. & KINALSKA, I. Soluble L-selectin levels in type I diabetes mellitus: a surrogate marker for disease activity?, 320
KRAMPS, T.A.N. *see* OEHEN, S.
KRANZER, K., BAUER, M., LIPFORD, G.B., HEEG, K., WAGNER, H. & LANG, R. CpG-oligodeoxynucleotides enhance T-cell receptor-triggered interferon-γ production and up-regulation of CD69 via induction of antigen-presenting cell-derived interferon type I and interleukin-12, 170
KREUTZ, W. *see* MÜLLER, B.
KRIEG, T. *see* YAMAMOTO, T.

LACY, P. *see* VELAZQUEZ, J.R.
LAMONT, A.G. *see* FRANCIS, J.N.
see also MACHT, L.M.
LANG, A.K. *see* SEARLE, R.F.
LANG, R. *see* KRANZER, K.
LANZA, F. *see* HIGGINBOTTOM, A.
LEE, M.-H., KATAOKA, T., HONJO, N., MAGAE, J. & NAGAI, K. *In vivo* rapid reduction of alloantigen-activated $CD8^+$ mature cytotoxic T cells by inhibitors of acidification of intracellular organelles, prodigiosin 25-C and concanamycin B, 243
LEHNER, T., WANG, Y., CRANAGE, M., TAO, L., MITCHELL, E., BRAVERY, C., DOYLE, C., PRATT, K., HALL, G., DENNIS, M., VILLINGER, F. & BERGMEIER, L. Up-regulation of β-chemokines and down-modulation of CCR5 co-receptors inhibit simian immunodeficiency virus in non-human primates, 569
LIN, T.-J. *see* DÉRY, R.E.
LIPFORD, G.B. *see* KRANZER, K.
LÓPEZ-MACÍAS, C. *see* OEHEN, S.
LYDYARD, P.M. *see* BODMAN-SMITH, M.D.

MACHT, L.M., ELSON, C.J., KIRWAN, J.R., GASTON, J.S.H., LAMONT, A.G., THOMPSON, J.M. & THOMPSON, S. J. Relationship between disease severity and responses by blood mononuclear cells from patients with rheumatoid arthritis to human heat-shock protein 60, 208
MADSEN, H.O. *see* JINQUAN, T.
MAEKAWA, Y. *see* SAKAI, T.
MAGAE, J. *see* LEE, M.-H.
MAILE, R., ELSEGOOD, K.A., HARDING, T.C., UNEY, J.B., STEWART, C.E.H., BANTING, G. & DAYAN, C. M. Effective formation of major histocompatibility complex class II–peptide complexes from endogenous antigen by thyroid epithelial cells, 367
MALLING, H.-J. *see* JINQUAN, T.
MARSHALL-CLARKE, S. *see* TASKER, L.
MATSUZAKI, G. *see* NINOMIYA, T.
MCCULLOUGH, B. *see* HIGGINBOTTOM, A.
MELCHER, A.A. *see* TODRYK, S.M.
MÉNARD, G. *see* DÉRY, R.E.
MERKENSCHLAGER, M. *see* BARCENAS-MORALES, G.

METCALFE, D.D. *see* NILSSON, G.
MIGITA, K. *see* HIDA, A.
MILNE, C.D. *see* DÉRY, R.E.
MITCHELL, E. *see* LEHNER, T.
MIYAGOE, Y. *see* IWAO, M.
MIYAZAKI, J. *see* SAKAI, T.
MONK, P.N. *see* HIGGINBOTTOM, A.
MOQBEL, R. *see* DÉRY, R.E.
see also VELAZQUEZ, J.R.
MORENO, A. *see* ALLENDE, L.M.
MORGAN, B P. *see* RUSHMERE, N.K.
MOZES, E. *see* BROSH, N.
MÜLLER, B., FISCHER, B. & KREUTZ, W. An acidic microenvironment impairs the generation of non-major histocompatability complex-restricted killer cells, 375

NAGAI, K. *see* LEE, M.-H.
NAKAMURA, H. *see* HIDA, A.
NAKANO, Y. *see* SAKAI, T.
NAKASHIMA, T. *see* HIDA, A.
NICOL, A., NIEDA, M., KOEZUKA, Y., PORCELLI, S., SUZUKI, K., TADOKORO, K., DURRANT, S. & JUJI, T. Human invariant $V\alpha24^+$ natural killer T cells activated by α-galactosylceramide (KRN7000) have cytotoxic anti-tumour activity through mechanisms distinct from T cells and natural killer cells, 229
NIEDA, M. *see* NICOL, A.
NILSSON, G., METCALFE, D.D. & TAUB, D. D. Demonstration that platelet-activating factor is capable of activating mast cells and inducing a chemotactic response, 314
NINOMIYA, T., TAKIMOTO, H., MATSUZAKI, G., HAMANO, S., YOSHIDA, H., YOSHIKAI, Y., KIMURA, G. & NOMOTO, K. $V\gamma1^+$ γδ T cells play protective roles at an early phase of murine cytomegalovirus infection through production of interferon-γ, 187
NISHIMURA, Y. *see* KONDO, A.
NITTA, Y. *see* SAKAI, T.
NOMOTO, K. *see* NINOMIYA, T.

O'BRIEN, D.P. *see* THATCHER, T.H.
OEHEN, S, JUNT, T., LÓPEZ-MACÍAS, C. & KRAMPS, T. A. N. Antiviral protection after DNA vaccination is short lived and not enhanced by CpG DNA, 163
OHNO, N. *see* KINGETSU, I.
OHNO, N., IDE, T., SAKAGUCHI, M., INOUYE, S. & SAITO, S. Common antigenicity between Japanese cedar (*Cryptomeria japonica*) pollen and Japanese cypress (*Chamaecyparis obtusa*) pollen, II. Determination of the cross-reacting T-cell epitope of Cry j 1 and Cha o 1 in mice, 630
OHSAWA, M. *see* YOSHINO, S.
OHUCHI, K. *see* SHIRAISHI, M.
OIKAWA, S. *see* SHIRAISHI, M.
OLIVÉ, T. *see* ALLENDE, L.M.
OVIEDO-ORTA, E., HOY, T. & EVANS, W. H. Intercellular communication in the immune system: differential expression of connexin40 and 43, and perturbation of gap junction channel functions in peripheral blood and tonsil human lymphocyte subpopulations, 578

PALMER, J. *see* SEARLE, R.F.
PARKHOUSE, R.M.E. *see* YANG, H.
PARR, E.L. *see* PARR, M.B.
PARR, M.B. & PARR, E. L. Interferon-γ up-regulates intercellular adhesion molecule-1 and vascular cell adhesion molecule-1 and recruits lymphocytes into the vagina of immune challenged mice challenged with herpes simplex virus-2, 540
PARTRIDGE, L.J. *see* HIGGINBOTTOM, A.
PODOLIN, P.L., WEBB, E.F., REDDY, M., TRUNEH, A. & GRISWOLD, D. E. Inhibition of contact sensitivity in human $CD4^+$ transgenic mice by human CD4-specific monoclonal antibodies: $CD4^+$ T-cell depletion is not required, 287
PORCELLI, S. *see* NICOL, A.
POULSEN, L.K. *see* JINQUAN, T.
PRATT, K. *see* LEHNER, T.

QUAN, S. *see* JINQUAN, T.

RADSAK, M. *see* WAGNER, C.
RAGNO, S. *see* TASCON, R.E.
REDDY, M. *see* PODOLIN, P.L.
RHODES, S.G., BUDDLE, B.M., HEWISON, R.G. & VORDERMEIER, H. M. Bovine tuberculosis: immune responses in the peripheral blood and at the site of active disease, 195
RIBBENS, C., DAYER, J.-M. & CHIZZOLINI, C. CD40–CD40 ligand (CD154) engagement is required but may not be sufficient for human T-helper 1 cell induction of interleukin-2- or interleukin-15-driven, contact-dependent, interleukin-1β production by monocytes, 279
RUSHMERE, N.K., VAN DEN BERG, C.W. & MORGAN, B. P. Production and functional characterization of a soluble recombinant form of mouse CD59, 326

SAHA, A. *see* BASU, A.
SAITO, S. *see* KINGETSU, I.
see also OHNO, N.; TODA, M.
SAKAGUCHI, M. *see* KINGETSU, I.
see also OHNO, N.; TODA, M.
SAKAI, H. *see* HIDA, A.
SAKAI, T., HISAEDA, H., NAKANO, Y., ISHIKAWA, H., MAEKAWA, Y., ISHII, K., NITTA, Y., MIYAZAKI, J. & HIMENO, K. Gene gun-mediated delivery of an interleukin-12 expression plasmid protects against infections with the intracellular protozoan parasites *Leishmania major* and *Trypanosoma cruzi* in mice, 615
SASATOMI, E. *see* YOSHINO, S.
SATO, H. *see* TODA, M.
SEARLE, R.F., BROMAGE, S.J., PALMER, J., CURRY, J.E. & LANG, A. K. Human amniotic fluid lacks interleukin-2 and interleukin-15 but can interact with the β-chain of the interleukin-2 receptor, 411
SHAPIRA, L. *see* HOURI-HADDAD, Y.
SHARMA, V., SPARKS, J.L. & VAIL, J. D. Human B-cell lines constitutively express and secrete interleukin-16, 266
SHEN, J. *see* BAO, S.
SHIRAISHI, M., HIRASAWA, N., OIKAWA, S., KOBAYASHI, Y. & OHUCHI, K. Analysis of histamine-producing cells at the late phase of allergic inflammation in rats, 600
SILVA, R.A., GOMES, M.S. & APPELBERG, R. Minor role played by type I tumour necrosis factor receptor in the control of *Mycobacterium avium* proliferation in infected mice, 203
SKOV, P.S. *see* JINQUAN, T.
SOARES, C.S. *see* TASCON, R.E.
SOSKOLNE, W.A. *see* HOURI-HADDAD, Y.
SPARKS, J.L. *see* SHARMA, V.
STAVROPOULOS, E. *see* TASCON, R.E.
STEWART, C.E.H. *see* MAILE, R.
STREILEIN, J.W. *see* KEZUKA, T.
see also KITAZAWA, T.
SUN, G.-H. *see* SUN, K.-H.
SUN, K.-H., YU, C.-L., TANG, S.-J. & SUN, G.-H. Monoclonal anti-double-stranded DNA autoantibody stimulates the expression and release of IL-1β, IL-6, IL-8, IL-10 and TNF-α from normal human mononuclear cells involving in the lupus pathogenesis, 352
SUZUKI, K. *see* NICOL, A.

TADOKORO, K. *see* NICOL, A.
TAKEBE, Y. *see* TODA, M.
TAKEDA, S. *see* IWAO, M.
TAKEMORI, T. *see* TODA, M.
TAKIMOTO, H. *see* NINOMIYA, T.
TANAKA, T. *see* KONDO, A.
TANG, S.-J. *see* SUN, K.-H.
TANIGUCHI, Y. *see* TODA, M.
TAO, L. *see* LEHNER, T.
TASCON, R.E., SOARES, C.S., RAGNO, S., STAVROPOULOS, E., HIRST, E.M.A. & COLSTON, M.J. *Mycobacterium tuberculosis*-activated dendritic cells induce protective immunity in mice, 473
TASKER, L. & MARSHALL-CLARKE, S. Antigen receptor signalling in apoptosis-resistant mutants of WEHI 231 cells, 385

TAUB, D.D. *see* NILSSON, G.
THATCHER, T.H., O'BRIEN, D.P., ALTUWAIJRI, S. & BARTH, R. K. Increasing the frequency of T-cell precursors specific for a cryptic epitope of hen-egg lysozyme converts it to an immunodominant epitope, 235
THOMPSON, J.M. *see* MACHT, L.M.
THOMPSON, S.J. *see* FRANCIS, J.N.
see also MACHT, L.M.
TODA, M., SATO, H., TAKEBE, Y., TANIGUCHI, Y., SAITO, S., INOUYE, S., TAKEMORI, T. & SAKAGUCHI, M. Inhibition of immunoglobulin E response to Japanese cedar pollen allergen (Cry j 1) in mice by DNA immunization: different outcomes dependent on the plasmid DNA inoculation method, 179
TODRYK, S.M., MELCHER, A.A., DALGLEISH, A.G. & VILE, R. G. Heat shock proteins refine the danger theory, 334
TRUNEH, A. *see* PODOLIN, P.L.
TSUJIKAWA, K. *see* IWAO, M.

UNEY, J.B. *see* MAILE, R.
URAYAMA, S. *see* HIDA, A.

VAIL, J.D. *see* SHARMA, V.
VAN DEN BERG, C.W. *see* RUSHMERE, N.K.
VARELA, P. *see* ALLENDE, L.M.
VELAZQUEZ, J.R., LACY, P. & MOQBEL, R. Replenishment of RANTES mRNA expression in activated eosinophils from atopic asthmatics, 591
VILE, R.G. *see* TODRYK, S.M.
VILLINGER, F. *see* LEHNER, T.
VORDERMEIER, H.M. *see* RHODES, S.G.
VUKUSIC, B. *see* WITHER, J.

WAGNER, C., BÜRGER, A., RADSAK, M., BLUM, S., HUG, F. & HÄNSCH, G. M. Fibronectin synthesis by activated T lymphocytes: up-regulation of a surface-associated isoform with signalling function, 532
WAGNER, H. *see* KRANZER, K.
WAHID, F. *see* BARCENAS-MORALES, G.
WANG, Y. *see* LEHNER, T.
WEBB, E.F. *see* PODOLIN, P.L.
WELKER, P., GRABBE, J., GIBBS, B., ZUBERBIER, T. & HENZ, B. M. Nerve growth factor-β induces mast-cell marker expression during *in vitro* culture of human umbilical cord blood cells, 418
WILKINSON, I. *see* HIGGINBOTTOM, A.
WITHER, J. & VUKUSIC, B. T-cell tolerance induction is normal in the (NZB × NZW)F_1 murine model of systemic lupus erythematosus, 345

YAGITA, H. *see* IWAO, M.
YAMAMOTO, H. *see* IWAO, M.
YAMAMOTO, T., HARTMANN, K., ECKES, B. & KRIEG, T. Mast cells enhance contraction of three-dimensional collagen lattices by fibroblasts by cell–cell interaction: role of stem cell factor/c-kit, 435
YAMASAKI, S. *see* HIDA, A.
YANG, H. & PARKHOUSE, R. M. E. Characterization of the porcine γδ T-cell receptor structure and cellular distribution by monoclonal antibody PPT27, 504
YOSHIDA, H. *see* NINOMIYA, T.
YOSHIDA, N. *see* AONO, A.
YOSHIKAI, Y. *see* NINOMIYA, T.
YOSHINO, S., SASATOMI, E. & OHSAWA, M. Bacterial lipopolysaccharide acts as an adjuvant to induce autoimmune arthritis in mice, 607
YOSHIZAKI, K. *see* AONO, A.
YOUINOU, P.Y. *see* BODMAN-SMITH, M.D.
YU, C.-L. *see* SUN, K.-H.

ZINGER, H. *see* BROSH, N.
ZUBERBIER, T. *see* WELKER, P.

Subject Index

Amniotic fluid:
lack of interleukin-2 and interleukin-15, 411
Anterior chamber-associated immune deviation:
evidence for multiple CD95–CD95 ligand interactions, 451
Antibodies, to cytokines:
in human immunoglobulin preparations, 113
Antigen presentation:
by thyroid epithelial cells, 367
Antigenicity:
of Japanese cedar pollen, 625
Apoptosis:
and antigen receptor signalling, 385
Apoptosis, of macrophages:
regulation by nuclear factor 'B and caspases, 553
Apoptosis, T cell:
and differential human leucocyte antigen class II expression, 249

B cells, bovine:
activation through immunoglobulin M and CD40, 272

C5a receptor expression:
in macrophages and renal proximal tubular, 38
CD4 engagement:
promotion of interleukin-4 synthesis, 394
CD9, human:
cross-linking with immunoglobulin E receptor, 546
CD18 genomic deletion:
in a leucocyte adhesion-deficiency patient, 440
CD40 ligand:
polymorphism, 54
CD59 mouse:
soluble recombinant, 326
CD59 rat:
production and functional analysis, 46
CD95 ligand:
in breast cancer, 69
CD98 and integrin signalling, 62
Connexins:
in human tonsillar lymphocytes, 578
Contact sensitivity:
in human $CD4^+$ transgenic mice, 287
CpG motif:
effects on dendritic cells, 361
CXC chemokine receptor 4 expression, on $CD4^+$ T cells:
regulation by interleukin-4 and interleukin-10, 402
Cytokine release, from lupus human mononuclear cells:
by anti-DNA, 352
Cytotoxic T cells, mature:
elimination by acidification inhibitors, 243

Dendritic cells:
activation by bacterial DNA sequences, 1
Dendritic cells, carcinogen modified:
induction of suppression, 16
Dendritic cells, costimulatory signals from:
after exposure to ultraviolet B, 296
Dendritic cells, *Mycobacterium tuberculosis* activated:
protective immunity, 473
Dendritic cells, splenic:
in murine leishmaniasis, 305
Dendritic cell vaccination:
anti-leukaemia responses, 8
Disaccharides:
regulation of T-cell adhesion, 87

DNA immunization:
immunotherapy for pollinosis, 179
DNA vaccination:
role of immune stimulatory sequences, 163

Fas–fas ligand interaction:
in graft-versus-host disease, 94
Fibronectin:
synthesis by activated T lymphocytes, 532

γδ T cells:
FcγRIII positive, 498
in cytomegalovirus infection, 187
γδ T cells, in skin:
effect of terminal deoxynucleotidyl transferase expression in fetal thymus, 489
γδ T-cell receptor, porcine:
characterization, structure and cellular distribution, 504

Heat shock proteins:
danger theory, 334
Heat shock protein 60, responses of mononuclear cells to:
in rheumatoid arthritis, 208
Heat shock protein 65 peptide, intranasal:
exacerbation of pristane-induced arthritis, 338
Histamine-producing cells:
at late phase of allergic inflammation, 600

$IgG2a^b$:
constitutive presentation of, 510
Immune response gene expression:
recessive, 221
Inducible nitric oxide synthase expression:
in infected trout, 153
Interferon-γ:
upregulation of intercellular adhesion molecule 1 and vascular cell adhesion molecule 1 in the vagina, 540
role in intestinal salmonella infection, 464
Interleukin-1β, monocyte:
induction by T helper 1 cells, 279
Interleukin-5, gene promoter:
DNAse footprinting, 101
Interleukin-10:
effect on antigen presentation of dendritic cells, 523
Interleukin-12 gene expression plasmid:
protection against intracellular parasites, 615
Interleukin-12 receptor β_1 and β_2, rat:
expression *in vivo*, 109
Interleukin-16:
human B cell, 266

Killer cells, non-major histocompatibility complex restricted:
inhibition by acidic pH, 375

Lectins, plant:
mucosal immunogenicity, 30
Lipopolysaccharide:
role in autoimmune arthritis, 607
L-selectin:
in type I diabetes, 320

Macrophage in T-cell interactions:
in bovine tuberculosis, 23
Mast cells:
enhancement of contraction of three-dimensional collagen lattices, 435

Merosin:
interaction with very late antigen-6 in the thymus, 481
Mls^+ B-cell activation, 147
Monocyte chemoattractant protein-3 gene expression:
in human endothelial cells, 561

Natural killer cells:
anti-tumour activity, 229
Nerve growth factor:
and cord-blood mast cells, 418

Platelet-activation factor:
activation of mast cells, 314
Porphyromonas gingivalis:
cytokine response to, 215
Prostaglandin, regulation:
from fetal membranes, 124

Regulated upon activation, normal T-cell expressed and secreted mRNA regulation:
in activated eosinophils, 591

Simian immunodeficiency virus transmission, in non-human parasites:
effect of β chemokines and CCR5 co-receptor, 569
Spα, mouse:
cloning and characterization, 78

Steroid hormones, endogenous:
role in T helper 2-mediated autoimmunity, 141

T helper 1 and T helper 2 profiles in *Haemonchus contortus*-infected sheep, 458
T-cell activation:
enhancement by CpG oligodeoxynucleotides, 170
T-cell epitopes, cross-reactivity:
in Japanese cedar pollen, 630
T-cell lines:
specific for complementarity determining region peptides, 257
T-cell precursors:
for cryptic epitope of hen-egg lysozyme, 235
Tolerance, T cell:
in NZB/NZW mice, 345
Tuberculosis, bovine:
immune responses, 195
Tumour necrosis factor production:
by macrophages and mast cells, 427
Tumour necrosis factor receptor type I:
role in *Mycobacterium avium* infection in mice, 203

Ultraviolet B radiation, effect on contact hypersensitivity:
role of interleukin-10, 134